

The ABCs of Your child's CPP journey

Your guide for taking important steps throughout their central precocious puberty (CPP) treatment journey

Activate the Prescription

Begin Treatment

Continue and Complete Treatment

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension) LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month, 11.25 mg and 30 mg for 3-month, and 45 mg for 6-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

It is not known if LUPRON DEPOT-PED is safe and effective in children less than 1 year old.

Safety Considerations Your child should not receive LUPRON DEPOT-PED if she/he is allergic to any of the ingredients or she is or may become pregnant. Increased signs and symptoms of puberty during the first 2 to 4 weeks of treatment may occur. Contact your child's doctor if signs of puberty continue after the second month of treatment.

Please see additional Important Safety Information on pages 12-15.

Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Now that you and your child's doctor have chosen LUPRON DEPOT-PED to treat your child's CPP, there are a few important things to know.

This brochure is designed to help guide you through and inform you about the treatment process and the support available through these ABCs:

Activate the Prescription

Getting your child's LUPRON DEPOT-PED prescription filled may take a few more steps than filling other prescriptions.

Begin Treatment

LUPRON DEPOT-PED is usually administered in your child's doctor's office as an injection. Your child's pediatric endocrinologist will decide on the appropriate starting dose for your child and will monitor your child's response to treatment.

Continue and Complete Treatment

Every child is unique, so it is important that you stay on track with appointments and for your child's treatment progress to be monitored regularly by their doctor.

Safety Considerations (cont'd) Some people taking gonadotropin-releasing hormone (GnRH) agonists, like LUPRON DEPOT-PED, have had new or worsened mental (psychiatric) problems. Mental (psychiatric) problems may include emotional symptoms such as crying, irritability, restlessness (impatience), anger, and acting aggressively. Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

Activate the Prescription



STEP 1: YOUR CHILD'S DOCTOR SUBMITS THE PRESCRIPTION

After you and your child's doctor have determined that LUPRON DEPOT-PED is right for your child, the doctor will send the prescription to a **specialty pharmacy** that handles medications that require special care.

The following may be required to process the prescription:

- Benefits Verification — the process of confirming your insurance coverage
- Prior Authorization — a special approval that some insurance plans require

Please see additional Important Safety Information on pages 12-15. Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)



STEP 2: YOUR INSURANCE PROVIDER REVIEWS THE PRESCRIPTION

If your insurance provider approves coverage, this means that your child will be able to start treatment!

If your insurance provider does not approve coverage, your child's doctor can file an appeal.



STEP 3: LUPRON DEPOT-PED ARRIVES

Once your child's LUPRON DEPOT-PED prescription is approved, it will be shipped directly from the specialty pharmacy to your child's doctor's office or to your home.

Contact your child's doctor's office right away to schedule the first injection appointment.

Safety Considerations (cont'd) Seizures have been observed in patients taking GnRH agonists, like LUPRON DEPOT-PED, and the risk of seizure may be higher in patients with or without a history of seizures, epilepsy, brain or brain vessel (cerebrovascular) problems or tumors, and in patients taking medications that have been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs). Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.



Please see additional Important Safety Information on pages 12-15. Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Begin Treatment

LUPRON DEPOT-PED is administered by injection in your child's doctor's office. Your child's dose will determine how often you need to make an appointment for additional injections.

LUPRON DEPOT-PED OFFERS FLEXIBLE & INDIVIDUALIZED DOSING OPTIONS

DOSING OPTIONS

6-month dosing	3-month dosing	1-month dosing
2 injections each year	4 injections each year	12 injections each year

Your child's pediatric endocrinologist decides on the appropriate starting dose. They will want to monitor your child a few times per year and adjust the dose as needed.

Scheduling options: If it's hard to get to appointments during regular office hours, tell your child's doctor. Some offices can work with your schedule.

Safety Considerations (cont'd) Increased pressure in the fluid around the brain can happen in children taking GnRH agonist medicines, including LUPRON DEPOT-PED. Call your child's doctor right away if your child has any of the following symptoms during treatment with LUPRON DEPOT-PED: headache, eye problems (including blurred vision, double vision, and decreased eyesight), eye pain, ringing in the ears, dizziness, or nausea. Your child's pubertal development could begin again if the injection schedule is not followed.

WHAT TO EXPECT

...AT THE DOCTOR'S OFFICE. Any appointment at the doctor's office may cause your child to be a little anxious, especially one in which they will get an injection.



SCAN THE QR CODE FOR TIPS TO HELP RELAX YOUR CHILD

www.lupronped.com/starting-therapy

...AFTER THE FIRST INJECTION, some pain and irritation at the injection site may occur. **Call your child's doctor** if these symptoms get worse, persist, or if severe reactions occur.



After the first few weeks of treatment, LUPRON DEPOT-PED can cause an increase in some hormones. During this time, you may notice more signs of puberty in your child, including vaginal bleeding in girls. **Don't worry!** This is normal. These signs should go away. If they continue beyond the second month, notify your child's doctor.

Please see additional Important Safety Information on pages 12-15. Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Continue and Complete Treatment

THINGS TO REMEMBER

- Keep your child's doctor appointments
- Follow the prescribed injection schedule
- Signs of puberty may begin again if injections are missed
- The doctor will do regular exams and blood tests to check for signs of puberty



MONITORING YOUR CHILD'S TREATMENT

Every child is unique, so how your child's signs and symptoms progress and how they respond to treatment may be different from other children. Your child's pediatric endocrinologist will check on your child's progress.

Since LUPRON DEPOT-PED offers different dosage choices, the doctor will be able to adjust the dosage based on your child's response to treatment.

Safety Considerations (cont'd) LUPRON DEPOT-PED should not be taken if your child is:

- Allergic to GnRH, GnRH agonist medicines, or any ingredients in LUPRON DEPOT-PED. See the end of the Medication Guide for a complete list of ingredients in LUPRON DEPOT-PED.
- Pregnant or becomes pregnant. LUPRON DEPOT-PED can cause birth defects or loss of the baby. If your child becomes pregnant, call your child's doctor.



Regular monitoring allows your child's pediatric endocrinologist to adjust the dose as necessary so that your child gets the amount of medicine that's right for them.

If you have any concerns about the effectiveness of your child's dosage level, be sure to bring them up with the doctor.

Sometimes the initial dosage needs to be adjusted—one way to know this is by **looking for signs and symptoms of puberty AFTER the second month of treatment.**



SCAN THE QR CODE TO LEARN WHAT TO LOOK OUT FOR

www.lupronped.com/starting-therapy

Call your child's doctor right away if your child experiences any new or worsened mental (psychiatric) symptoms or problems or if your child has a seizure while taking LUPRON DEPOT-PED.

When your child reaches the age when puberty should occur, the doctor will stop your child's treatment to allow their puberty to start back up again.

Please see additional Important Safety Information on pages 12-15. Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Support PLUS:

WE'RE HERE FOR YOU.
EVERY STEP OF THE WAY.

Support PLUS can help you with:



Understanding Insurance Coverage

A Support PLUS Insurance Specialist can talk you through your coverage, explain what your out-of-pocket costs may be, and help identify potential savings options for your child's LUPRON DEPOT-PED prescription.



Live Nurse Support Hotline*

You can speak with our certified nurses about CPP and treatment with LUPRON DEPOT-PED anytime over the course of your child's treatment—at no cost to you.

**Call the toll-free hotline:
1-855-LUPRON-P (1-855-587-7667)**

Support PLUS Service	Days of Operation	Hours of Operation
Nurse Hotline (Nurse On-Call)	Monday - Friday	10 AM–7 PM CT
Insurance Coverage Assistance (HUB)	Monday - Friday	7 AM–7 PM CT

*Certified nurses are provided by AbbVie and do not work under the direction of your healthcare professional (HCP) or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

ELIGIBLE COMMERCIALLY INSURED PATIENTS† MAY PAY AS LITTLE AS \$10 PER PRESCRIPTION OF LUPRON DEPOT-PED



**Scan the QR code to see if you are eligible
and request an Instant Savings Card**

www.lupronped.com/lupron-support-plus#savings-card

LUPRON DEPOT-PED Co-pay Full Terms and Conditions

†Terms and Conditions apply. This benefit covers LUPRON DEPOT-PED® (leuprolide acetate for depot suspension). Eligibility: Available to patients with commercial insurance coverage for LUPRON DEPOT-PED who meet eligibility criteria. Co-pay assistance program is not available to patients receiving reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare (including Part D), Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law or by the patient's health insurance provider. If at any time a patient begins receiving drug coverage under any such federal, state, or government-funded healthcare program, patient will no longer be able to use the LUPRON DEPOT-PED Instant Savings Card and patient must call LUPRON DEPOT-PED at 1-855-587-7667 to stop participation. Patients residing in or receiving treatment in certain states may not be eligible. Patients may not seek reimbursement for value received from the LUPRON DEPOT-PED Instant Savings Card program from any third-party payers. Offer subject to change or discontinuation without notice. Restrictions, including monthly maximums, may apply. Subject to all other terms and conditions, the maximum annual benefit that may be available solely for the patient's benefit under the co-pay assistance program is \$4,000 per calendar year for patients receiving LUPRON DEPOT-PED every month or \$7,000 per calendar year for patients receiving LUPRON DEPOT-PED every 3 and/or 6 months. The actual application and use of the benefit available under the co-pay assistance program may vary on a monthly, quarterly, and/or annual basis depending on each individual patient's plan of insurance and other prescription drug costs. This assistance offer is not health insurance. By redeeming this card, you acknowledge that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer. To learn about AbbVie's privacy practices and your privacy choices, visit <https://privacy.abbvie>

**Please see additional Important Safety Information on pages 12-15.
Please see accompanying Consumer Brief Summary
or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf>
for full Prescribing Information.**

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month, 11.25 mg and 30 mg for 3-month, and 45 mg for 6-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

It is not known if LUPRON DEPOT-PED is safe and effective in children less than 1 year old.

Important Safety Information for LUPRON DEPOT-PED

What is the most important information I should know about LUPRON DEPOT-PED?

- **During the first 2 to 4 weeks of treatment, LUPRON DEPOT-PED can cause an increase in some hormones.** During this time, you may notice more signs of puberty in your child, including vaginal bleeding. **Call your child's doctor if these signs continue after the second month of treatment with LUPRON DEPOT-PED.**
- **Some people taking gonadotropin-releasing hormone (GnRH) agonists like LUPRON DEPOT-PED have had new or worsened mental (psychiatric) problems.** Mental (psychiatric) problems may include emotional symptoms such as:
 - Crying
 - Irritability
 - Restlessness (impatience)
 - Anger
 - Acting aggressive

Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

- **Some people taking GnRH agonists like LUPRON DEPOT-PED have had seizures. The risk of seizures may be higher in people who:**
 - Have a history of seizures
 - Have a history of epilepsy
 - Have a history of brain or brain vessel (cerebrovascular) problems or tumors
 - Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)

Seizures have also happened in people who have not had any of these problems. **Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.**

- **Increased pressure in the fluid around the brain can happen in children taking gonadotropin-releasing hormone (GnRH) agonist medicines, including LUPRON DEPOT-PED. Call your child's doctor right away if your child has any of the following symptoms during treatment with LUPRON DEPOT-PED:**

- Headache
- Eye pain
- Eye problems including blurred vision, double vision, and decreased eyesight
- Ringing in the ears
- Dizziness
- Nausea

LUPRON DEPOT-PED should not be taken if your child is:

- Allergic to GnRH, GnRH agonist medicines, or any ingredients in LUPRON DEPOT-PED. See the end of the Medication Guide for a complete list of ingredients in LUPRON DEPOT-PED.
- Pregnant or becomes pregnant. LUPRON DEPOT-PED can cause birth defects or loss of the baby. If your child becomes pregnant, call your child's doctor.

Before your child receives LUPRON DEPOT-PED, tell their doctor about all of your child's medical conditions, including if they:

- Have a history of mental (psychiatric) problems
- Have a history of seizures
- Have a history of epilepsy
- Have a history of brain or brain vessel (cerebrovascular) problems or tumors
- Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)
- Are breastfeeding or plan to breastfeed. It is not known if LUPRON DEPOT-PED passes into the breast milk

Important Safety Information for LUPRON DEPOT-PED (cont'd)

Tell your child's doctor about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will your child receive LUPRON DEPOT-PED?

- Your child's doctor should do tests to make sure your child has CPP before treating them with LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is given as a single-dose injection into your child's muscle each month, every 3 months, or every 6 months by a doctor or trained nurse. Your child's doctor will decide how often your child will receive the injection.
- Keep all scheduled visits to the doctor. If a scheduled dose is missed, your child may start having signs of puberty again. The doctor will do regular exams and blood tests to check for signs of puberty.

What are the possible side effects of LUPRON DEPOT-PED?

LUPRON DEPOT-PED may cause serious side effects. See "What is the most important information I should know about LUPRON DEPOT-PED?"

- **The most common side effects of LUPRON DEPOT-PED received 1 time each month include:**
 - Injection site reactions such as pain, swelling, and abscess
 - Weight gain
 - Pain throughout body
 - Headache
 - Acne or red, itchy rash and white scales (seborrhea)
 - Serious skin rash (erythema multiforme)
 - Mood changes
 - Swelling of vagina (vaginitis), vaginal bleeding, and vaginal discharge

- **The most common side effects of LUPRON DEPOT-PED received every 3 months include:**

- Injection site reactions such as pain and swelling
- Weight gain
- Headache
- Mood changes

- **The most common side effects of LUPRON DEPOT-PED received every 6 months include:**

- Injection site reactions such as pain, swelling, and abscess
- Headache
- Fever
- Weight gain
- Mood changes
- Itching
- Fracture
- Upper Stomach pain
- Pain in extremities
- Breast tenderness
- Diarrhea
- Rash
- Difficulty sleeping
- Bleeding
- Back pain
- Chest pain
- Nausea and vomiting
- Ligament sprain
- Excessive sweating

These are not all the possible side effects of LUPRON DEPOT-PED. **Call your child's doctor for medical advice about side effects.**

This is the most important information to know about LUPRON DEPOT-PED. For more information, talk to your child's doctor or healthcare provider.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Please see accompanying Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.



HERE FOR YOU EVERY STEP OF THE WAY

Scan the QR code to learn more about Support PLUS

<https://www.lupronped.com/lupron-support-plus>



**Contact Support PLUS, toll-free, at
1-855-LUPRON-P (1-855-587-7667),
Monday through Friday, 7 AM–7 PM CT**

**Please see pages 12-15 for Use and Important Safety Information.
Please see accompanying Consumer Brief Summary
or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf>
for full Prescribing Information.**

abbvie

LUPRON DEPOT-PED[®]
(leuprolide acetate for depot suspension)

© 2023 AbbVie. All rights reserved.
LUPRON DEPOT-PED[®] and its design are
registered trademarks of AbbVie Inc.
US-LUPR-230055 May 2023